



# Review: Long-term effect of perioperative smoking cessation programmes

Hanne Tønnesen & Thordis Thomsen

## Abstract

**Background** Preoperative smoking cessation programmes have been reviewed recently regarding the immediate effect on postoperative complications and smoking habits. The objective of this review was therefore to evaluate long-term effects of perioperative smoking cessation programmes.

**Methods** PubMed, Cochrane, Embase, CINAHL databases were searched for randomised clinical trials on perioperative smoking cessation intervention programmes that included follow-up for smoking. The literature was evaluated and data were extracted from the included papers. The review involved meta-analyses.

**Results** The 10 included RCTs were presented in 12 papers and communications, involving 1,369 patients. Only 5 RCTs had a follow up of 12 months. The RR in the perioperative period was 1.90 (95% CI: 1.65 to 2.18), and after 12 months 1.57 (1.09 to 2.26). The subgroup analyses of the intensive 6-8 weeks programmes revealed a high RR of 5.89 (3.49 to 9.93).

**Conclusion** Smoking cessation intervention programmes were effective on short and long-term. The intensive programmes of six to eight weeks duration seemed to be most effective.

## About the AUTHORS

### Hanne Tønnesen

WHO-Collaborating Centre for Evidence-Based Health Promotion in Hospitals and Health Services, Bispebjerg University Hospital

### Thordis Thomsen

Dept Anaesthesiology and Intensive Medicine, Herlev Hospital, Copenhagen

### Contact:

Hanne Tønnesen  
hanne.tonnesen@bbh.regionh.dk

## Introduction

The association between smoking and postoperative complications is well established (10;11) and more than three hundred papers have been published since dr. Morton first described the increased risk of pulmonary problems in smokers compared to non-smokers in 1944 (12). However, the first randomised clinical trial (RCT) on smoking cessation in relation to surgery was published more than fifty years later (11;13). It was soon followed by eight more RCTs and further ongoing studies are to be published in the coming years, all together gathering substantial knowledge on the effect of different smoking cessation programmes in the perioperative period (14).

Preoperative smoking cessation programmes have been reviewed recently regarding the immediate effect on postoperative complications and smoking habits (11;14-16). However, a possible long-term effect and related prognostic factors still needed to be investigated further. The aim of this review was therefore to evaluate long-term effects of perioperative smoking cessation programmes.

## Material

The inclusion criteria were RCTs on perioperative smoking cessation interventions among smokers undergoing surgery and postoperative follow up for smoking cessation. The programme should involve personal contact; it could be brief or intensive intervention, with or without pharmacotherapy. The control group could receive treatment as usual or placebo.


In total, 10 trials were identified and presented in 12 papers including 4 papers on long-term follow-up on original trials (2;4;29;30). Two trials did not distinguish between smoking reduction and smoking cessation in the outcome data, and were therefore excluded (27;28). This review included 10 trials for further evaluation; the trial profile is given in figure 1.

The main outcome measurement was smoking cessation up to one year after the intervention, either continuous or point prevalence.



## Research and Best Practice

**Figure 1** Trial profile for the systematic review



### Methods

A systematic literature search was performed in the databases PubMed, Embase, CINAHL and the Cochrane Library; supplemented by hand search. There was no time or language restriction. The search strategy included participants (smoking OR tobacco use disorder) AND intervention OR smoking cessation OR tobacco use cessation OR smoking intervention OR tobacco use intervention OR smoking counselling OR tobacco use counselling OR patient education OR preoperative care OR perioperative care OR preoperative preventive care OR perioperative preventive care OR health promotion program OR preoperative health promotion program OR perioperative health promotion program) AND relation to surgery (surgery OR operation OR surgical procedure OR perioperative intervention OR postoperative complication\* OR intraoperative complication\*).

The search included RCTs as well as clinical controlled trials (CCT), and reviews with or without meta-analysis in order not to overlook weakly defined RCTs; the only limit was 'all adults'.

The quality of the studies was evaluated through the Cochrane Collaboration's tool for assessing risk of bias (35).

Data from the 10 RCTs were extracted regarding number of participants, follow-ups and drop outs, types of intervention and control programmes, quit rates and validation as well as regarding risk of bias;

sequence generation, allocation concealment, blinding, incomplete data, selective reporting and other bias.

Mantel-Haenszel methods were used to calculate risk ratios (RRs) and corresponding 95 per cent confidence intervals (CI). RRs were calculated using available case analysis (35).

Metaanalyses were performed using the fixed effect method. Heterogeneity among studies was calculated using the  $I^2$  statistic describing the percentage of the variability in effect estimates, which is due to heterogeneity rather than sampling error. Metaanalyses were performed only if the  $I^2$  of heterogeneity was below 40%.

Subgroup analyses were conducted according to the intensity of the smoking cessation programmes; brief intervention (BI) and intensive intervention (II) in 4-8 weeks programmes, with and without nicotine replacement therapy (NRT), bupropion and varenicline; the setting (intervention with and without relation to the surgical setting); and study quality (high and low). Review Manager (RevMan) version 5.0 (The Nordic Cochrane Centre, The Cochrane Collaboration, Copenhagen, Denmark) was used for data analysis. The results were presented as risk rates (RR) and 95% confidence interval (95% CI) (35).

### Results

The 10 included trials involved 1,369 patients from Denmark, Sweden, United Kingdom, Australia, Canada and the US. The characteristics are described in table 1.

The risk of bias was relatively low, because all trials reported adequate sequence generation and allocation concealment, and they were free from selective reporting or other bias. All trials reported intention to treat analyses. However, four trials were unclear about the blinding procedure (3;4;29;36), two did not clearly address incomplete outcome data (36;37), and two reported point abstinence from smoking instead of continuous abstinence (38;39).

Four trials included 12 months follow-up, three trials for 3 and 6 months respectively.

The analyses showed significant quit rates on short term preoperatively and immediately postoperatively in 7 of 9 studies (Figure 2, a). However, the quit rates reduced over time with the exception of the studies that tested intensive interventions (figure 2 b,c,d,e).

Only the quit rates of the 6-8 weeks intensive inter-




Research and Best Practice

**Table 1** Characteristics of the included 10 trials presented in 12 papers/communications.

References	Preop. smoking cessation program	Duration (Weeks)	Validation	No. of patients	No. of drop-outs	Follow-ups (Months post)
Thomsen et al. 2010(2)	BI + NRT	1-2	Yes	130	17	Peri 1+3+6+12
Sadr Azodi et al. 2009(29) +Lindström et al. 2008(40)	II + NRT + hotline	4	Yes	117	19 15	Peri 1+12
Nåsell et al. 2010(1)	II + NRT + hotline	8	Yes	105	11	Peri 1½
Villebro et al. 2008(4) + Møller et al. 2002(13)	II + NRT	6-8	Yes	120	19 12	Peri 1+12
Sørensen et al. 2007(36)	BI (telephone) + NRT	1	Yes	180	31	Peri 3
Andrew et al. 2006(3)	BI (letter)	4	No	102	1	Peri
Warner 2005(47)	NRT	0	No	121	5	1+6
Wolfender 2005(39)	BI (computer and telephone) + NRT	1-2	No	210	29	Peri 3
Myles 2004(37)	Bupropion + BI (telephone)	7	Yes	47	23	Peri
Ratner 2004(30)	BI	1-3	Yes	237	69	Peri 6+12

BI = brief intervention. II = intensive intervention. NRT = nicotine replacement treatment. Peri = perioperative. Post = postoperative

**Figure 2** Results of preoperative smoking cessation programmes on quit rates in the perioperative period (a), and at follow up after 1 (b), 3 (c), 6 (d), and 12 (e) months. The blue colour reflects the intensive programmes.



vention programmes stayed significant during the total follow-up period (figure 2). Accordingly, the subgroup analyses of the intensive programmes revealed a high RR of 2.87 (1.50 to 5.58) for long-term smoking cessation (figure 3b).

**Discussion**


We found a significant effect of the perioperative smoking cessation programmes on short-term, while there was no clear overall effect on longer term. However, the intensive intervention programmes of 6-8weeks showed a significantly increased smoking cessation rate after one year (4;13;29;40). No studies followed up on the quit rate for more than one year, and they did not include long-term functionality and health, mortality or costs over time.

Studies of smoking intervention in other groups of hospital patients have recently been reviewed by dr. Rigotti and her colleagues (41). They found that only programmes lasting for one month or more after discharge were effective, thus sustaining our results. Nevertheless, these



## Research and Best Practice

**Figure 3** Sub-group analysis of the results of intensive smoking cessation programmes on quit rates in the perioperative period (a) and at follow-up after 12 months (b).



findings do not exclude that minimal smoking cessation programmes might be useful for some sub-groups of hospitalised smokers or smokers in other settings. The intensive smoking intervention programmes also have a clear beneficial effect on the development of complications after surgery in smokers compared to the brief intervention programmes (14), and they seem most attractive for implementation in surgical settings.

However, the bias and limitations of our review should be kept in mind. Especially, the relatively small sample sizes of several of the included studies may be followed by a significant risk of type 2-failure, thereby overlooking minor effects. On the other hand, the use of point prevalence and self-reported quit rates without validation may overestimate an outcome (38;39). Furthermore, the patient groups differed regarding primary diagnosis and co-morbidity, smoking history and other factors influencing the effect of smoking cessation programmes. According to the consecutiveness of included patients the heterogeneity of the materials may be high. On the other hand the consecutiveness would improve the possibility of generalising and translation of the results. All studies were performed in high-income Western countries, and care should be taken if translated to other countries, cultures and patient groups.

From a clinical perspective, the intensive smoking cessation intervention programmes of 4-8 weeks of duration are preferable for surgical patients due to the immediate risk reduction of postoperative complications previously described as well as due to the beneficial long term effect on the quit rate. Implementation through evidence-based clinical guidelines and follow-up for effect could easily be established using the HPH Model for Documentation of Health Promotion activities (42) and a clinical qual-

ity registry such as the Danish national smoking cessation database ([www.scdb.dk](http://www.scdb.dk)). These tools allow ongoing exchange of knowledge and experience as well as identification and updating of the most effective programmes for the benefit of the individual patient and the society as a whole.

The patients are positive, when it comes to smoking cessation programs in the perioperative period, especially the intensive programmes of 4-8 weeks duration are requested (4;40;43;44). This is in contrast to the fear of stigmatising smokers, when recommending smoking cessation before surgery, which was hypothesised previously (45;46).

From a scientific point of view, future studies should be powered to include follow-up after longer time, e.g. 3-5 years, for smoking habits as well as for long-term postoperative complications and functionality, mortality and costs.

In conclusion, this review supports that the briefer perioperative smoking cessation intervention programmes are effective on short-term only, while the intensive programmes of six to eight weeks duration are effective on long-term as well.

**Competing interests:** None declared.

### Contributors

Conception and design: HT, TT

Acquisition of data: HT, TT

Analysis and interpretation of data: TT

Drafting the article: HT

Revising the article critically for important intellectual content: TT

Approving the article: HT, TT

### Reference List

- (1) Nåsell H, Adami J, Samnegård E, Tønnesen H, Ponzer S. Effect of Smoking Cessation Intervention on Results of Acute Fracture Surgery - A Randomized Controlled Trial. *J Bone Joint Surg (Am)* 2010;92:1335-42.
- (2) Thomsen T, Tønnesen H, Okholm M, Kroman N, Maibom A, Sauerberg M. Brief smoking cessation intervention in relation to breast cancer surgery: A randomized controlled trial. *Nicotine Tob Res.* 2010;12:1118-24
- (3) Andrews K, Bale P, Chu J, Cramer A, Aveyard P. A randomized controlled trial to assess the effectiveness of a letter from a consultant surgeon in causing smokers to stop smoking pre-operatively. *Public Health* 2006 April;120:356-8.
- (4) Villebro N, Pedersen T, Møller AM, Tønnesen H. Long-term effects of a preoperative smoking cessation programme. *CRJ* 2008;2:175-82.



## Research and Best Practice

- (5) Wolfenden L, Wiggers J, Campbell E, Knight J. Pilot of a preoperative smoking cessation intervention incorporating post-discharge support from a Quitline. *Health Promot J Austr* 2008;19:158-60.
- (6) Greenhalgh RM, Belch JJ, Brown LC, Gaines PA, Gao L, Reise JA, Thompson SG. The adjunct benefit of angioplasty in patients with mild to moderate intermittent claudication (MIMIC) managed by supervised exercise, smoking cessation advice and best medical therapy: results from two randomised trials for stenotic femoropopliteal and aortoiliac arterial disease. *Eur J Vasc Endovasc Surg* 2008;36:680-8.
- (7) Wallstrom M, Nilsson F, Hirsch JM. A randomized, double-blind, placebo-controlled clinical evaluation of a nicotine sublingual tablet in smoking cessation. *Addiction* 2000;95:1161-71.
- (8) Sørensen LT, Jørgensen LN, Zillmer R, Vange J, Hemmingsen U, Gottrup F. Transdermal nicotine patch enhances type I collagen synthesis in abstinent smokers. *Wound Repair Regen* 2006;14:247-51.
- (9) Sørensen LT, Karlsmark T, Gottrup F. Abstinence from smoking reduces incisional wound infection: a randomized controlled trial. *Ann Surg.* 2003;238:1-5.
- (10) Møller AM, Tønnesen H. Risk reduction: perioperative smoking intervention. *Best Pract Res Clin Anaesthesiol* 2006;20:237-48.
- (11) Warner DO. Perioperative abstinence from cigarettes: physiologic and clinical consequences. *Anesthesiology* 2006;104:356-67.
- (12) Morton HJV. Tobacco smoking and pulmonary complications after operation. *Lancet* 1944;1:368-70.
- (13) Møller AM, Villebro N, Pedersen T, Tønnesen H. Effect of preoperative smoking intervention on postoperative complications: a randomised clinical trial. *Lancet* 2002;359:114-7.
- (14) Thomsen T, Tønnesen H, Møller AM. Effect of preoperative smoking cessation interventions on postoperative complications and smoking cessation. *Br J Surg* 2009;96:451-61.
- (15) Thomsen T, Villebro N, Moller AM. Intervention for preoperative smoking cessation (review). *Cochrane Database Syst Rev* 2010 (7).
- (16) Tønnesen H, Nielsen PR, Lauritzen JB, Møller AM. Smoking and alcohol intervention before surgery: evidence for best practice. *Br J Anaesth* 2009;102:297-306.
- (17) Motovska Z, Widimsky P, Petr R, Bilkova D, Marinov I, Simek S, Kala P. Factors influencing clopidogrel efficacy in patients with stable coronary artery disease undergoing elective percutaneous coronary intervention: statin's advantage and the smoking "paradox". *J Cardiovasc Pharmacol* 2009;53:368-72.
- (18) Wilson-MacDonald J, Fairbank J, Frost H, Yu LM, Barker K, Collins R, Campbell H. The MRC spine stabilization trial: surgical methods, outcomes, costs, and complications of surgical stabilization. *Spine (Phila Pa 1976)* 2008;33:2334-40.
- (19) Hogue CW, Jr., Fucetola R, Hershey T, Nassief A, Birge S, Vila-Roman VG, Barzilai B, Thomas B, Schechtman KB, Freedland K. The role of postoperative neurocognitive dysfunction on quality of life for postmenopausal women 6 months after cardiac surgery. *Anesth Analg* 2008;107:21-8.
- (20) Albertal M, Cura F, Escudero AG, Thierer J, Trivi M, Padilla LT, Belardi J. Mechanism involved in the paradoxical effects of active smoking following primary angioplasty: a subanalysis of the protection of distal embolization in high-risk patients with acute myocardial infarction trial. *J Cardiovasc Med (Hagerstown)* 2008;9:810-2.
- (21) Pluss CE, Karlsson MR, Wallen NH, Billing E, Held C. Effects of an expanded cardiac rehabilitation programme in patients treated for an acute myocardial infarction or a coronary artery by-pass graft operation. *Clin Rehabil* 2008;22:306-18.
- (22) Dastoor SF, Travan S, Neiva RF, Rayburn LA, Giannobile WV, Wang HL. Effect of adjunctive systemic azithromycin with periodontal surgery in the treatment of chronic periodontitis in smokers: a pilot study. *J Periodontol* 2007;78:1887-96.
- (23) Weisz G, Cox DA, Garcia E, Tchong JE, Griffin JJ, Guagliumi G, Stuckey TD, Rutherford BD, Mehran R, Aymong E, Lansky A, Grines CL, Stone GW. Impact of smoking status on outcomes of primary coronary intervention for acute myocardial infarction--the smoker's paradox revisited. *Am Heart J* 2005;150:358-64.
- (24) Taira DA, Seto TB, Ho KK, Krumholz HM, Cutlip DE, Berezin R, Kuntz RE, Cohen DJ. Impact of smoking on health-related quality of life after percutaneous coronary revascularization. *Circulation* 2000;102:1369-74.
- (25) Glasgow RE, Estabrooks PA, Marcus AC, Smith TL, Gaglio B, Levinson AH, Tong S. Evaluating initial reach and robustness of a practical randomized trial of smoking reduction. *Health Psychol* 2008;27:780-8.
- (26) Glasgow RE, Gaglio B, Estabrooks PA, Marcus AC, Ritzwoller DP, Smith TL, Levinson AH, Sukhanova A, O'Donnell C, Ferro EF, France EK. Long-term results of a smoking reduction program. *Med Care* 2009;47:115-20.
- (27) Hughes JA, Sanders LD, Dunne JA, Tarpey J, Vickers MD. Reducing smoking. The effect of suggestion during general anaesthesia on postoperative smoking habits. *Anaesthesia* 1994;49:126-8.
- (28) Sørensen LT, Jørgensen T. Short-term pre-operative smoking cessation intervention does not affect postoperative complications in colorectal surgery: a randomized clinical trial. *Colorectal Dis* 2003;5:347-52.
- (29) Sadr Azodi O, Lindstrom D, Adami J, Tonnesen H, Nasell H, Gilljam H, Wladis A. The efficacy of a smoking cessation programme in patients undergoing elective surgery: a randomised clinical trial. *Anaesthesia* 2009;64:259-65.
- (30) Ratner PA, Johnson JL, Richardson CG, Botorff JL, Moffat B, Mackay M, Fofonoff D, Kingsbury K, Miller C, Budz B. Efficacy of a smoking-cessation intervention for elective-surgical patients. *Res Nurs Health* 2004;27:148-61.
- (31) Møller AM, Pedersen T, Villebro N, Nørgaard P. Impact of lifestyle on perioperative smoking cessation and postoperative complication rate. *Prev Med* 2003;36:704-9.
- (32) Simon JA, Solkowitz SN, Carmody TP, Browner WS. Smoking cessation after surgery. A randomized trial. *Arch Intern Med* 1997;157:1371-6.
- (33) Stanislaw AE, Wewers ME. A smoking cessation intervention with hospitalized surgical cancer patients: a pilot study. *Cancer Nurs* 1994;17:81-6.
- (34) Lindeman CA. Nursing intervention with the presurgical patient: effectiveness and efficiency of group and individual preoperative teaching. *Nurs Res* 1972;21:196-209.
- (35) Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.0. 2008.
- (36) Sørensen LT, Hemmingsen U, Jørgensen T. Strategies of smoking cessation intervention before hernia surgery--effect on perioperative smoking behavior. *Hernia* 2007;11:327-33.
- (37) Myles PS, Leslie K, Angliss M, Mezzavia P, Lee L. Effectiveness of bupropion as an aid to stopping smoking before elective surgery: a randomised controlled trial. *Anaesthesia* 2004;59:1053-8.
- (38) Andrews DW, Silverman CL, Glass J, Downes B, Riley RJ, Corn BW, Werner-Wasik M, Curran WJ, Jr., McCune CE, Rosenwasser RH, . Preservation of cranial nerve function after treatment of acoustic neuromas with fractionated stereotactic radiotherapy. Preliminary observations in 26 patients. *Stereotact Funct Neurosurg* 1995;64:165-82.
- (39) Wolfenden L, Wiggers J, Knight J, Campbell E, Rissel C, Kerridge R, Spigelman AD, Moore K. A programme for reducing smoking in pre-operative surgical patients: randomised controlled trial. *Anaesthesia* 2005;60:172-9.
- (40) Lindström D, Azodi OS, Wladis A, Tønnesen H, Linder S, Nasell H, Ponzer S, Adami J. Effects of a perioperative smoking cessation intervention on postoperative complications: a randomized trial. *Ann Surg* 2008;248:739-45.
- (41) Rigotti NA, Munafo MR, Stead LF. Smoking cessation interventions for hospitalized smokers: a systematic review. *Arch Intern Med* 2008;168:1950-60.
- (42) Tønnesen H, Christensen ME, Groene O, O'Riordan A, Simonelli F, Suurorg L, Morris D, Vibe P, Himel S, Hansen PE. An evaluation of a model for the systematic documentation of hospital based health promotion activities: results from a multicentre study. *BMC Health Serv Res* 2007;7:145.
- (43) Møller AM, Villebro NM. Preoperative smoking intervention: What do patients think? A qualitative study. *Ugeskr Laeger* 2004;166:3714-8.
- (44) Thomsen T, Esbensen BA, Samuelsen S, Tønnesen H, Møller AM. Brief pre-operative smoking cessation counselling in relation to breast cancer surgery: A qualitative study. *Eur J Oncol Nurs* 2009;13:344-9.
- (45) Glantz L. Should smokers be refused surgery? *BMJ* 2007;334:21.
- (46) Peters MJ. Should smokers be refused surgery? *BMJ* 2007;334:20.
- (47) Warner DO, Patten CA, Ames SC, Offord KP, Schroeder DR. Effect of nicotine replacement therapy on stress and smoking behavior in surgical patients. *Anesthesiology* 2005;102:1138-46.